



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

	1	5	9	
Third	2 3	6 7	10 11	. . . .
	4	8	12	

All the combinations involving 1, 2, 3, 4 only have been used, and thereby 1 has played *with* each 2, 3, 4 once and twice against each. Since the players are identically involved, *mutatis mutandis*, all the other players will have a similar experience.

361. Proposed by C. E. GITHENS, Ph. D., Wheeling, W. Va.

Find three integral values for  $[-10+9\sqrt{-3}]^{\frac{1}{3}}+[-10-9\sqrt{-3}]^{\frac{1}{3}}$ . A solution not involving a cubic is desired.

I. Solution by the PROPOSER.

I. Put  $(-10+9\sqrt{-3})^{\frac{1}{3}}+(-10-9\sqrt{-3})^{\frac{1}{3}}=\frac{1}{2}[(-80+72\sqrt{-3})^{\frac{1}{3}}+(-80-72\sqrt{-3})^{\frac{1}{3}}]$ .

Let  $(-80+72\sqrt{-3})^{\frac{1}{3}}=\sqrt[3]{x}+\sqrt[3]{y}$  and  $(-80-72\sqrt{-3})^{\frac{1}{3}}=\sqrt[3]{x}-\sqrt[3]{y}$ .

Then,  $[6400-(-15552)]^{\frac{1}{3}}=x-y=28$ , and, therefore,

$(-80+72\sqrt{-3})^{\frac{1}{3}}=\sqrt[3]{x+28}+\sqrt[3]{x}$ . Hence, by raising both sides to the third power,  $(4x+28)\sqrt[3]{x+28}+(4x+84)\sqrt[3]{x}=-80+72\sqrt{-3}$ .

Put  $4x+28\sqrt[3]{x+28}=-80$ , and  $4x+84\sqrt[3]{x}=72\sqrt{-3}$ .

Let  $4x+28=-80$  or factor of  $-80$ ;  $-40$ ,  $-20$ ,  $-16$ ,  $-10$ ,  $-9$ ,  $-8$ ,  $-5$ ,  $-4$ ,  $-2$ , and

$4x+84=72\sqrt{-3}$  or factor of  $72$ ;  $36$ ,  $24$ ,  $18$ ,  $12$ ,  $9$ ,  $8$ ,  $6$ ,  $4$ ,  $3$ ,  $2$ .

Subtracting,  $-56=-20-(36)$ .

$\therefore 4x+28=-20$  and  $x=-12$ .

[1]  $\frac{1}{2}[(-80+72\sqrt{-3})^{\frac{1}{3}}+(-80-72\sqrt{-3})^{\frac{1}{3}}]$   
 $=\frac{1}{2}[(\sqrt[3]{x+28}+\sqrt[3]{x})+(\sqrt[3]{x+28}-\sqrt[3]{x})]=4$ , answer.

II. Similarly with  $(-80+72\sqrt{-3})^{\frac{1}{3}}=\sqrt[3]{x_0+28}-\sqrt[3]{x_0}$ , in which the factors  $-80$  and  $24$  added to eliminate the  $4x$ 's produces

$x_0=-27$  and  $\frac{1}{2}[(\sqrt[3]{x_0+28}-\sqrt[3]{x_0})+\sqrt[3]{x_0+28}+\sqrt[3]{x_0}]=-1$ , answer.

III. With  $-\sqrt[3]{x_1+28}+\sqrt[3]{x_1}$  and  $-16$  and  $72$  as factors as above,  $x=-3$  and  $\frac{1}{2}[-\sqrt[3]{x_1+28}+\sqrt[3]{x_1}+(-\sqrt[3]{x_1+28}-\sqrt[3]{x_1})]^{\frac{1}{3}}=-5$ , answer.

IV. The root  $-\sqrt[3]{x_2+28}-x_2$  produces no two factors of  $-80$  and  $72$  whose difference or sum equals  $\pm 56$ ; hence it is not a root, which is as it should be, for the numerical equation is an example of the "irreducible case" in the Cardan solution of a cubic whose equation is  $x^3-21x+20=0$ .

II. Solution by J. SCHEFFER, A. M., Hagerstown, Maryland.

Putting  $(-10+9\sqrt{3}\sqrt{-1})=\rho(\cos\phi+\sin\phi\sqrt{-1})$ , we get  $\tan\phi=-\frac{9}{10}\sqrt{3}$ .  $\rho=\sqrt{343}$ , and  $(-10+9\sqrt{3}\sqrt{-1})^{\frac{1}{3}}+(-10-9\sqrt{3}\sqrt{-1})^{\frac{1}{3}}=2\sqrt{7}\cos\frac{1}{3}\phi$ .

$$\therefore 4\cos^3\frac{1}{3}\phi-3\cos\phi+\frac{1}{4}\sqrt{7}=0.$$

By trial,  $\cos\frac{1}{3}\phi=\frac{2}{7}\sqrt{7}$ , and dividing the last trinomial by  $\cos\frac{1}{3}\phi-\frac{2}{7}\sqrt{7}$ , we get  $4\cos^2\frac{1}{3}\phi+\frac{8}{7}\sqrt{7}\cos\frac{1}{3}\phi-\frac{5}{7}=0$ ; whence  $\cos\frac{1}{3}\phi=-\frac{5}{14}\sqrt{7}$ ,  $\cos\frac{1}{3}\phi=\frac{1}{14}\sqrt{7}$ .

$\therefore$  The three values required are 4, -5, 1.

Also solved by A. M. Harding.

## GEOMETRY.

386. Proposed by DANIEL KRETH, Oxford, Iowa.

Construct the triangle, having given, the vertical angle, the sum of the three sides, and the perpendicular.

I. Solution by H. PRIME, Boston, Massachusetts.

Let  $ABC$  be the required triangle,  $C$  the given angle. On  $AB$  produced take  $BE=BC$ . On  $BA$  produced take  $AF=AC$ . Let  $O$  be the center of circle  $ECF$ . Then we have the angles  $FOE=2(BEC+AFC)=ABC+BAC=\text{supplement of } C$ .


Hence, to construct the triangle, on  $EF=\text{the given sum of the three sides}$  form the isosceles triangle  $EOF$ , making  $EOF=\text{the supplement of the given vertex angle}$  (or  $EOF=OFE=\text{one half the given angle}$ ). About  $O$  as center describe the arc  $EF$ . Parallel to  $EF$  and at a distance from it equal to the given altitude draw a line meeting the arc at  $C$  and  $C'$ . Draw  $CA$  and  $CB$ , making the angles  $ACF=AEC$  and  $BCE=BEC$ .  $ABC$  is the required triangle.

II. Solution by C. N. SCHMALL, New York City, and A. M. HARDING, University of Arkansas.

Construct an angle  $A$  equal to the given vertical angle. Lay off  $AD$  and  $AE$  each equal to *half* the given sum of the sides. Describe a circle touching these lines in  $D$  and  $E$ . With  $A$  as center and radius equal to the given perpendicular, describe a circle  $MN$ . By a well known method draw a line tangent to *both* these circles touching in  $R$  and  $S$ , respectively, and cutting the sides in  $B$  and  $C$ . Then  $ABC$  is the triangle required.

Proof.  $BR=BD$ ,  $CR=CE$ .

$\therefore BC=BD+CE$ ; hence the triangle has the given perimeter. Also,  $AS$  is perpendicular to  $BC$ ; therefore the triangle has the required altitude. Q. E. D.



Also solved by J. Scheffer and A. H. Holmes.